

# A Review Study: Effect of Growth Factors on Human Mesenchymal Stem Cells Differentiation into Cartilage Tissue

Babak Pourmollaabbasi<sup>1</sup>, Batool Hashemibeni<sup>2\*</sup>, Ebrahim Esfandiari<sup>2</sup>

1. Department of Tissue Engineering, School of Basic Engineering, Islamic Azad University of Najafabad, Isfahan, Iran.

2. Department of Anatomical Sciences, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran.



Babal Pourmollaabbasi Graduated from MSc tissue engineering, departement of tissue engineering at school of basic engineering in Islamic azad university of Najafabad, Isfahan, Iran. His interests are study and working on stem cells and scaffold for regeneration and tissue engineering of cartilage and bone defects.

Article info:

Received: 17 Jun. 2015

Accepted: 09 Aug. 2015

## Key Words:

Mesenchymal stem cell,  
Growth factor, Chondro  
cartilage, Tissue engineering

## ABSTRACT

Hyaline cartilage is a vascular and neural tissue with scanty chondrocytes and limited regenerative ability. After some serious injuries of the cartilage, healing process will take place through the formation of fibrocartilage structures. Currently, tissue engineering and cell therapy are 2 interesting therapeutic fields dealing with regenerative medicine. In this regard, tissue regeneration has found mesenchymal stem cells (MSCs) with self-renewal and multipotential abilities as the best candidates for this process. Growth and differentiation of MSCs are induced by growth factors. The purpose of this review article is to evaluate the effect of growth factors and their signaling pathways involved in differentiation of mesenchymal stem cells into chondrocytes in vitro conditions.

## 1. Introduction

One of the most crippling diseases is arthralgia which leads to people's disabilities [1]. Arthralgia is caused by different factors such as osteoarthritis or trauma resulting from destruction of hyaline cartilage. About 36 million American people with arthralgia are suffering from osteoarthritis or joint injuries resulting from sports activities [2].

Physicians, surgeons, and researchers have tried to treat and improve degenerated cartilage. Current treatments have resulted in insufficient repair of cartilage and instead fibrocartilage formation. The main reason for the partial cure is the lack of blood vessels and low mitosis of chondrocytes in the joint cartilage. Thus, hyaline cartilage has a limited ability to repair its serious injuries [3, 4]. Successful efforts have been made over the past few years to cure joint cartilages by using mesenchymal stem cells (MSCs). The cells can be isolated from different sources such as bone marrow, adipose tissues, synovial

### \* Corresponding Author:

Batool Hashemibeni, PhD

Address: Department of Anatomical Sciences, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran.

Tel: +98 (31) 37929152

E-mail: hashemibeni@med.mui.ac.ir

membrane, dental pulp, umbilical cord blood, placental, and so on [5-10]. MSCs are undifferentiated cells with self-renewal and multipotential abilities [11,12]. They express specific surface markers like MSCA-1, STRO-1, CD271, CD105, CD90, CD73, CD44, and CD29 [13,14,15]. These cells can differentiate into other cells by induction of appropriate growth factors. The growth factors like transforming growth factor-beta (TGF- $\beta$ ), fibroblast growth factor (FGF), and insulin-like growth factor (IGF) have been applied to differentiate MSCs into chondrocytes [16-18]. This review article tries to evaluate the effect of growth factors and pathways of signaling involved in the differentiation of MSCs into chondrocytes.

## 2. Transforming growth factor-beta superfamily

Some studies have been carried out to determine the types of growth factors and their effects on growth, differentiation, and apoptosis of chondrocytes, ultimately, leading to the discovery of the superfamily of the transforming growth factor-beta (TGF- $\beta$ ) [19]. This family includes more than 35 members, and it is involved in different biological processes such as cell proliferation, differentiation, development of embryo, immunity reaction, inflammation, and repair [20,21].

TGF- $\beta$  family members affect in 2 separate signaling pathways, one is TGF/Nodal/Activin branch, and the other is bone morphogenic protein (BMP) signaling branch [22]. Recent studies have shown that the TGF- $\beta$  has 3 isotopes;  $\beta$ 1,  $\beta$ 2, and  $\beta$ 3 produced as inactive form until the peptide, depending on cloning, is separated from them Latency Associated Peptide (LAP) [23]. The transforming growth factor-beta (TGF- $\beta$ ) members bind to the serine/threonine kinase receptors (type I, II) on the cell surface. After binding TGF- $\beta$  members to their receptors, they get phosphorylated and activated [24]. Activated molecules trigger intracellular signaling via SMAD proteins (SMA=small body size, MAD=Mothers against decapentaplegic) [25]. The SMAD protein molecules that initiate the activity are Smad-2 and Smad-3 known as R-Smad.

### Three groups of SMAD protein molecules family

The family of SMAD protein molecules is divided into 3 groups:

- Receptor-regulated SMADs (R-SMAD), which include SMAD1, SMAD2, SMAD3, SMAD5 and SMAD8/9 [26],

- Common-mediator SMAD (co-SMAD), which includes only Smad4, interacting with R-SMADs to participate in signaling [27].

- Antagonistic or inhibitory SMADs (I-SMAD), which includes SMAD6 and SMMAD7, blocking the activation of R-SMADs and co-SMADs [28].

Also, the expression of SMAD 1/4/5 in chondrocytes causes upregulation of interleukin-1 (IL1) [29]. R-SMADs activation is related to phosphorylation of SXs region located at C-terminal, which has a different behavior relative to Activin receptor-Like Kinases (ALK) [30].

Recent studies have shown that Protein Phosphatase PPM1A dephosphorylates Smad 2/3 controls the signaling pathway of TGF- $\beta$  by phosphorylation and dephosphorylation of SMAD2 and 3 [31, 32]. Activated R-SMAD forms a complex with SMAD4 that is transferred into the nucleus and regulates the genes of chondrogenic differentiation [33]. Following the activation of the receptors, the mitogen-activated protein kinase (MAPK) may get activated. In fact, 3 pathways, including extracellular signal-regulated kinases (ERK), C-Jun-NH2-terminal Kinase, and P38 are considered for MPAKs.

The other members of SMADs named SMAD6 and 7 confine the activities of SMAD2 and SMAD3 [34] and inhibit the chondrogenesis of MSCs [35]. RUNX2 is a transcription factor involved in osteochondrogenic differentiation pathway. Study has shown that blocking of RUNX2 expression causes inhibition of osteogenesis in mice [36]. When SMAD6 and SMAD7 interact with RUNX2 in SMAD pathways, the rate of proteases increases, and these enzymes by destroying Runx2, prevent differentiation process [37]. R-SMADs can also be inhibited by other pathways, including SARA (SMAD-anchored for receptor activation) membranous proteins [38]. The most important members of TGF- $\beta$  superfamily that can be used for cartilage tissue engineering include TGF- $\beta$ 1, TGF- $\beta$ 2, BMP2, BMP4, BMP7, and GDF5 [39].

### The transforming growth factor-beta 1

It has been revealed that the transforming growth factor-beta 1 (TGF- $\beta$ 1) stimulates the production of extracellular matrix in chondrocytes [40]. Moreover, this growth factor inhibits the catabolic activity of IL-1 in cartilage tissue [41]. In vitro studies have shown that TGF- $\beta$ 1 affects the differentiation of MSCs into the chondrocytes. Additionally, TGF- $\beta$ 1 inhibits the production of type-I

collagen and increases the expression of type-II collagen and aggrecan genes in chondrogenic differentiation of MSCs [42].

### The transforming growth factor-beta 3

Hashemibeni et al. have found that TGF- $\beta$  promotes chondrogenesis of adipose-tissue-derived stem cells (ADSCs) in 3D scaffold-free pellet culture system [43]. Studies have shown that transforming growth factor-beta 3 (TGF- $\beta$ 3) enhances the synthesis of glycosaminoglycans (GAGs) in chondrogenesis of MSCs [44,45]. The findings showed that type II collagen, GAGs, and aggrecan are expressed in differentiated ADSCs in chondrogenic medium supplemented with TGF- $\beta$ 3. Meanwhile, they produce more AGC, GAG, and type I collagen in comparison to natural chondrocytes [46].

Esfandiari et al. have found that using both TGF $\beta$ 3 and physical (EF) inducers has the best outcomes in chondrogenesis, expression of SOX9, and type II collagen genes [47].

### Bone morphogenic proteins

Bone morphogenetic proteins (BMPs) are members of the transforming growth factor-beta superfamily that advance their signaling pathway through different pathways relative to TGF- $\beta$ . They have 4 receptors at the cell surface, including BMPRI $\alpha$  (ALK2, ALK3), BMPRI $\beta$  (ALK6), and BMPRII each of which activates different signaling pathways [48]. The BMP signaling pathway is initiated by binding ligand to its receptors and causing the activation of SMADs 1/5/8 by phosphorylation. Following the formation of complex by SMADs1/5/8 with SMAD4 and entering in nucleus, this complex stimulates the cascade of reacting and specific gene expression [49,50]. SMAD7 and suppressors of cytokine signaling (SOCS) have negative feedback effect on BMP signaling pathways [51].

### Bone morphogenic protein-2

Bone morphogenetic protein-2 (BMP2) increases extracellular matrix production (ECM) and reduces type-I collagen expression. Sekiya et al. have found that using BMP2 on MSCs increases the production of aggrecan due to the application of other members of TGF- $\beta$  [52].

### Bone morphogenic protein-4

Bone morphogenic protein 4 (BMP4) is an important and effective protein in osteogenesis and chondrogenesis. The results of studies conducted on the effects of

BMP4 on MSCs for differentiation into chondrocyte indicated that this growth factor leads to the production of chondroprogenitor lines and chondrogenic differentiation. In addition, BMP4 increases type-II collagen and aggrecan expression and at the same time, suppresses type-I and X collagens [53].

### Bone morphogenic protein-7

This growth factor is synthesized by the chondrocytes inducing the anabolic activity to support the damaged cartilage [54]. Studies indicated that BMP7 reduces MSCs proliferation, but it stimulates the cartilage extracellular matrix production. Cheng et al. have found that the applications of BMP7 in combination with TGF- $\beta$ 1 and IGF-I would result in the best effects in chondrogenesis [55].

### Growth differentiation factor

Researchers have found that among 15 members of subfamily of growth differentiation factors (GDFs), only GDF5 has chondrogenic potential and stimulates type-II collagen and aggrecan production in micromass or pellet culture systems [56, 57]. Also, GDF5 expresses in skeleton and joints development [58].

### Insulin-like growth factor-1

Insulin-like growth factor-1 (IGF1) plays the most important role in cartilage tissue. It causes homeostasis of cartilages and proteoglycan synthesis, proliferation, and differentiation of chondrocytes in the growth plates in vitro and in vivo [59, 60]. This growth factor repairs cartilage by increasing the matrix and type II collagen synthesis [61, 62]. The studies indicated that IGFI could induce differentiation of MSCs into cartilage, but when used with other growth factors such as TGF- $\beta$ 1 and BMP7, it exhibits better results [63].

The activity of insulin growth factors in tissues and blood circulation is regulated by a group of IGF-related proteins (IGFBP) so that some of them stimulate and others suppress the effects of IGFI. Studies have shown that IGFBP1, IGFBP2, IGFBP4, and IGFBP6 act as inhibitors while IGFBP3 serves as a stimulator in IGFI signaling pathway [64, 65].

The IGFI activity starts by binding to tyrosine kinase receptor (IGF-IR) followed by the activation of SHC (SH2-containing inositol phosphatase) and members of insulin receptor-substrate (IRS) family, including IRS1 and IRS2. Following the phosphorylation of IRS, 3 cas-

cade pathways of PI3K (phosphoinositol 3-kinase), ERK (extracellular signal regulated kinase), and MAPK (mitogen-activated protein kinase) are activated.

Activated P13K stimulates AKt and P70-S6 kinase and affects the cell survival and synthesis of the proteins. On the other hand, binding of Grb2 (Growth factor receptor-bound protein-2) to IGF-R causes the activation of IRS-1, SHC, Ras and Raf/MEK/ERK/MAPK. Research indicated that IGF1 stimulates proteoglycan synthesis in serum and synovial liquid. Studies have shown that IGF1 activates PI3K leading to chondrogenesis [66].

When chondrocytes are stimulated by insulin-like growth factor-1 (IGF1), PI3K causes proliferation of cells, gene expression, protein synthesis, type-II collagen and proteoglycans production, and prevention of apoptosis [67, 68]. Other studies indicated that IGF1 inhibits extracellular matrix production in the cartilage, thus IGF1 has opposite effects on cartilage tissues [69].

### **Fibroblast growth factor family**

A total of 23 members of fibroblast growth factor (FGF) family from FGF1 to FGF23 are known so far [70]. Fibroblast growth factors are heparin-associated polypeptides involved in neovascularization in vivo and in the growth of new blood vessels during the wound healing and embryogenesis.

The fibroblast growth factors in vitro conditions cause proliferation, differentiation, and cell migration leading to the production of protease in endothelial cells involved with tyrosine kinase receptors (FGFRs) and glucosaminoglycan (GAG) binding to heparin (HLGC) [71]. After the connection of fibroblast growth factor (FGF) to its receptor on the cell surface, the FGF activates the signaling pathways comprising of phosphoinositol kinase (PI3K) [72], phospholipase, mitogen-activated protein kinase (MAPK), extracellular signal-regulated kinases (ErK/p42/44), and P38 [73, 74]. Evidently, 2 members of the family of fibroblast growth factors; FGF-2 and FGF-18 (FGF), play the most important role in chondrogenic induction [75].

### **Fibroblast growth factor-2**

Fibroblast growth factor (FGF) is known as an important factor for the regulation of cartilage homeostasis, and its mitogenic role [76]. Using fibroblast growth factor-2 (FGF2) in monolayer culture prepares the required conditions for extracellular matrix synthesis and expression of specific phenotype in chondrocytes [77]. Stud-

ies have shown that FGF2 increases proliferation and production of proteoglycan in MSCs and also it induces human adipose-derived stem cells chondrogenesis in Transwell culture, which may be beneficial to cartilage tissue engineering [78].

In addition, FGF2 preserves the multipotency of the MSCs in vitro [79]. FGF2 in the joint cartilage and meniscus decreases the ratio of type-I collagen to type-II collagen and thus fibrocartilage replaces with hyaline cartilage for the repair of damaged tissue [80].

### **Fibroblast growth factor-18**

Fibroblast growth factor-18 (FGF18) induces the cell growth, tissue repair, tumor growth, homeostasis protection, and extracellular matrix (ECM) of cartilage formation [81]. In vitro studies demonstrated that when FGF18 binds to FGFR3, receptors suppress cell proliferation and improve the differentiation and production of ECM [82]. Tonia et al. showed that FGF18 has a positive effect on the chondrogenesis and can repair the damaged cartilage in animal model study [83].

## **3. Discussion**

Today, joint damages and pains are the most important problems that have affected human health in spite of life modernization. Current treatments such as subchondral drilling, arthroplasty, and autologous chondrocyte implantation do not have satisfactory results, as cartilage repair is incomplete. Thus, researchers have focused on the application of MSCs to repair damaged cartilage.

Using MSCs for desired cartilage tissue engineering requires appropriate growth factors like FGF, IGF, and TGF families. Over the past several years, in vitro and in vivo studies have been conducted to introduce MSCs into joint cartilage. Now, it is time to study the signaling pathways of each growth factor and determine what factors stimulate or inhibit the signaling pathway of chondrogenesis. With regard to the studies and research conducted so far, some positive effects and achievements have been attained, but occasionally some irreversible damages and negative effects have also been observed [84].

In summary, more research should be carried out on chondrogenic induction of MSCs and clinical application of engineered cartilage tissue.

## Acknowledgements

We would like to express our deep gratitude to Dr. Gohariyan, member of Novin darman garan company, for their service and generous support to conduct this research.

## References

- [1] Buckwalter JA, Mankin HJ. Articular cartilage Part II: degeneration and osteoarthritis, repair, regeneration, and transplantation. *Journal of Bone & Joint Surgery*. 1997; 79(4):612-32.
- [2] Temenoff JS, Mikos AG. Tissue engineering for regeneration of articular cartilage. *Biomaterials*. 2000; 21(5):431-440.
- [3] Stockwell RA. *Biology of cartilage cells*. Cambridge: Cambridge University Press; 1979.
- [4] Brown TD, Johnson RC, Saltzman CL, Marsh JL, Buckwalter JA. Post traumatic osteoarthritis: A first estimate of incidence, prevalence, and burden of disease. *Journal of Orthopaedic Trauma*. 2006; 20(10):739-744.
- [5] Winter A, Breit S, Parsch D, Benz K, Steck E, Hauner H, et al. Cartilage-like gene expression in differentiated human stem cell spheroids: A comparison of bone marrow-derived and adipose tissue-derived stromal cells. *Arthritis and Rheumatism*. 2003; 48(2):418-29.
- [6] Sakaguchi Y, Sekiya I, Yagishita K, Muneta T. Comparison of human stem cells derived from various mesenchymal tissues: Superiority of synovium as a cell source. *Arthritis & Rheumatism*. 2005; 52(8):2521-9.
- [7] Esfandiari E, Roshankhah S, Mardani M, Hashemibeni B, Naghsh E, Kazemi M, et al. The effect of high frequency electric field on enhancement of chondrogenesis in human adipose-derived stem cells. *Iranian Journal of Basic Medical Sciences*. 2014; 17(8):571-576.
- [8] Hashemibeni B, Razavi S, Esfandiari E, Karbasi S, Mardani M, Nadali F, et al. Designing of Human Cartilage Tissue, by differentiation of adipose-derived stem cells with BMP-6 in alginate scaffold. *Journal of Iranian Anatomical Sciences*. 2010; 8(31):117-27.
- [9] Kurth T, Hedbom E, Shintani N, Sugimoto M, Chen FH, Haspl M, et al. Chondrogenic potential of human synovial mesenchymal stem cells in alginate. *Osteoarthritis and Cartilage*. 2007; 15(10):1178-89.
- [10] Danišovič L, Varga I, Polák Š. Growth factors and chondrogenic differentiation of mesenchymal stem cells. *Tissue and Cell*. 2012; 44(2):69-73.
- [11] Bernardo ME, Emons JA, Karperien M, Nauta AJ, Willemze R, Roelofs H, et al. Human mesenchymal stem cells derived from bone marrow display a better chondrogenic differentiation compared with other sources. *Connective Tissue Research*. 2007; 48(3):132-40.
- [12] Kim HJ, Im GI. Chondrogenic differentiation of adipose tissue-derived mesenchymal stem cells: Greater doses of growth factor are necessary. *Journal of Orthopaedic Research*. 2009; 27(5):612-9.
- [13] Dominici ML, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini FC, Krause DS, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. *International Society for Cellular Therapy Position Statement*. 2006; 8(4):315-7.
- [14] Alipour R, Sadeghi F, Hashemibeni B, Zarkesh-Esfahani SH, Heydari F, Mousavi SB, et al. Phenotypic characterizations and comparison of adult dental stem cells with adipose-derived stem cells. *International Journal of Preventive Medicine*. 2010; 1(3):164-71.
- [15] Shafaei H, Esmaeili A, Mardani M, Razavi S, Hashemibeni B, Nasr-Esfahani MH, et al. Effects of human placental serum on proliferation and morphology of human adipose tissue-derived stem cells. *Bone Marrow Transplantation*. 2011; 46(11):1464-71.
- [16] Shanmugarajan TS, ByungSooKim H. Growth factors and signaling pathways in the chondrogenic differentiation of mesenchymal stem cells. *Tissue Engineering and Regeneration Medicine*. 2011; 8(3):292-9.
- [17] Kabiri A, Esfandiari E, Hashemibeni B, Kazemi M, Mardani M, Esmaeili A. Effects of FGF-2 on human adipose tissue derived adult stem cells morphology and chondrogenesis enhancement in Transwell culture. *Biochemical and Biophysical Research Communications*. 2012; 424(2):234-238.
- [18] Hashemibeni B, Goharian V, Esfandiari E, Sadeghi F, Fasihi F, Alipour R, et al. An animal model study for repair of tracheal defects with autologous stem cells and differentiated chondrocytes from adipose-derived stem cells. *Journal of Pediatric Surgery*. 2012; 47(11):1997-2003.
- [19] Davidson EB, Van Der Kraan PM, Van Den Berg WB. TGF- $\beta$  and osteoarthritis. *Osteoarthritis and Cartilage*. 2007; 15(6):597-604.
- [20] Gordon KJ, Blobel GC. Role of transforming growth factor- $\beta$  superfamily signaling pathways in human disease. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*. 2008; 1782(4):197-228.
- [21] Razavi S, Esfandiari E, Karbasi S, Mardani M, Sadeghi F, Esfahani MN, et al. Effect of TGF- $\beta$ 3 and BMP-6 growth factors on chondrogenic differentiation of adipose-derived stem cells in alginate scaffold. *Journal of Isfahan University of Medical Sciences*. 2010; 28(112):608-20.
- [22] Stewart R, Stojkovic M, Lako M. Mechanisms of self-renewal in human embryonic stem cells. *European Journal of Cancer*. 2006; 42(9):1257-72.
- [23] Hyytiäinen M, Penttinen C, Keski-Oja J. Latent TGF- $\beta$  binding proteins: Extracellular matrix association and roles in TGF- $\beta$  activation. *Critical Reviews in Clinical Laboratory Sciences*. 2004; 41(3):233-64.
- [24] Bertolino P, Deckers M, Lebrin F, ten Dijke P. TGF- $\beta$  signal transduction in angiogenesis and vascular disorders. *Chest*. 2005; 128(6):585-590.
- [25] Savage C, Das P, Finelli AL, Townsend SR, Sun CY, Baird SE, Padgett RW. *Caenorhabditis elegans* gene sma-2, sma-3, and sma-4 define a conserved family of transforming

- growth factor beta pathway components. Proceedings of the National Academy of Sciences of the United States of America. 1996; 93(2):790-794.
- [26] Wu JW, Hu M, Chai J, Seoane J, Huse M, Li C, et al. Crystal structure of a phosphorylated Smad2: Recognition of phosphoserine by the MH2 domain and insights on smad function in TGF- $\beta$  signaling. *Molecular Cell*. 2001; 8(6):1277-89.
- [27] Shi Y, Hata A, Lo RS, Massagué J, Pavletich NP. A structural basis for mutational inactivation of the tumour suppressor Smad4. *Nature*. 1997; 388(6637):87-93.
- [28] Itoh F, Asao H, Sugamura K, Heldin CH, ten Dijke P, Itoh S. Promoting bone morphogenetic protein signaling through negative regulation of inhibitory Smads. *EMBO Journal*. 2001; 20(15):4132-42.
- [29] Bau B, Haag J, Schmid E, Kaiser M, Gebhard PM, Aigner T. Bone morphogenetic protein-mediating receptor-associated Smads as well as common Smad are expressed in human articular chondrocytes but not up-regulated or down-regulated in osteoarthritic cartilage. *Journal of Bone and Mineral Research*. 2002; 17(12):2141-50.
- [30] Wrighton KH, Lin X, Feng XH. Termination of TGF- $\beta$  superfamily signaling. *Cell Research*. 2009; 19(1):8-20.
- [31] Lin X, Chen Y, Meng A, Feng X. Termination of TGF- $\beta$  superfamily signaling through SMAD dephosphorylation: A functional genomic view. *Journal of Genetics and Genomics*. 2007; 34(1):1-9.
- [32] Lin X, Duan X, Liang YY, Su Y, Wrighton KH, Long J, et al. PPM1A functions as a Smad phosphatase to terminate TGF $\beta$  signaling. *Cell*. 2006; 125(5):915-28.
- [33] Song B, Estrada KD, Lyons KM. Smad signaling in skeletal development and regeneration. *Cytokine and Growth Factor Reviews*. 2009; 20(5):379-88.
- [34] Konrad L, Scheiber JA, Bergmann M, Eickelberg O, Hofmann R. Identification of a new human Smad6 splice variant. *Andrologia*. 2008; 40(6):358-63.
- [35] Iwai T, Murai J, Yoshikawa H, Tsumaki N. Smad7 inhibits chondrocyte differentiation at multiple steps during endochondral bone formation and down-regulates p38 MAPK pathways. *Journal of Biological Chemistry*. 2008; 283(40):27154-64.
- [36] Hecht J, Seitz V, Urban M, Wagner F, Robinson PN, Stiege A, et al. Detection of novel skeletogenesis target genes by comprehensive analysis of a Runx2-/- mouse model. *Gene Expression Patterns*. 2007; 7(1):102-12.
- [37] Shen R, Chen M, Wang YJ, Kaneki H, Xing L, O'Keefe RJ, et al. Smad6 interacts with Runx2 and mediates Smad ubiquitin regulatory factor 1-induced Runx2 degradation. *Journal of Biological Chemistry*. 2006; 281(6):3569-76.
- [38] Hanyu A, Ishidou Y, Ebisawa T, Shimanuki T, Imamura T, Miyazono K. The N domain of Smad7 is essential for specific inhibition of transforming growth factor- $\beta$  signaling. *Journal of Cell Biology*. 2001; 155(6):1017-28.
- [39] Hatakeyama Y, Tuan RS, Shum L. Distinct functions of BMP4 and GDF5 in the regulation of chondrogenesis. *Journal of Cellular Biochemistry*. 2004; 91(6):1204-17.
- [40] Danisovic L, Lesny P, Havlas V, Teyssler P, Syrova Z, Kopani M, et al. Chondrogenic differentiation of human bone marrow and adipose tissue-derived mesenchymal stem cells. *Journal of Applied Biomedicine*. 2007; 5(4):139-50.
- [41] Awad HA, Halvorsen YC, Gimble JM, Guilak F. Effects of Transforming Growth Factor  $\beta$ 1 and Dexamethasone on the Growth and Chondrogenic Differentiation of Adipose-Derived Stromal Cells. *Tissue Engineering*. 2004; 9(6):1301-1312.
- [42] Abnaof K, Mallela N, Walenda G, Meurer SK, Seré K, Lin Q, et al. TGF- $\beta$  stimulation in human and murine cells reveals commonly affected biological processes and pathways at transcription level. *BMC Systems Biology*. 2014; 8(1):55.
- [43] Hashemibeni B, Razavi S, Esfandiari E, Karbasi S, Mardani M, Nasresfahani M. Induction of chondrogenic differentiation of human adipose-derived stem cells with TGF-R3 in pellet culture system. *Iran Journal of Basic Medical Sciences*. 2008; 11(1):10-17.
- [44] Yun-Feng RU, Lin DU, You WA, Yang WA, Pauline Poyee LU, Ting-ting TA, et al. Bone morphogenetic protein 2 promotes transforming growth factor  $\beta$ 3-induced chondrogenesis of human osteoarthritic synovium-derived stem cells. *Chinese Medical Journal*. 2010; 123(21):3040-8.
- [45] Ansar MM, Esfandiarii E, Mardani M, Hashemibeni B, Zarkesh-Esfahani SH, Hatf M, et al. A comparative study of aggrecan synthesis between natural articular chondrocytes and differentiated chondrocytes from adipose derived stem cells in 3D culture. *Advanced Biomedical Research*. 2012; 1(1):24.
- [46] Mardani M, Hashemibeni B, Ansar M, ZarkeshEsfahani H, Kazemi M, Goharian V, et al. Comparison between chondrogenic markers of differentiated chondrocytes from adipose derived stem cells and articular chondrocytes In vitro. *Iranian Journal of Basic Medical Sciences*. 2013; 16(6):763-773.
- [47] Hashemibeni B, Esfandiari E, Sadeghi F, Heidary F, Roshankhah S, Mardani M, Goharian V. An animal model study for bone repair with encapsulated differentiated osteoblasts from adipose-derived stem cells in alginate. *Iranian Journal of Basic Medical Sciences*. 2014; 17(11):854-859.
- [48] Nishimura R, Hata K, Matsubara T, Wakabayashi M, Yoneda T. Regulation of bone and cartilage development by network between BMP signaling and transcription factors. *Journal of Biochemistry*. 2012; 151(3):247-54.
- [49] Ying QL, Nichols J, Chambers I, Smith A. BMP induction of Id proteins suppresses differentiation and sustains embryonic stem cell self-renewal in collaboration with STAT3. *Cell*. 2003; 115(3):281-92.
- [50] Massagué J, Wotton D. Transcriptional control by the TGF- $\beta$ /Smad signaling system. *EMBO Journal*. 2000; 19(8):1745-54.
- [51] Rao M. Conserved and divergent paths that regulate self-renewal in mouse and human embryonic stem cells. *Developmental Biology*. 2004; 275(2):269-86.
- [52] Sekiya I, Larson BL, Vuoristo JT, Reger RL, Prockop DJ. Comparison of effect of BMP-2 and-6 on invitro cartilage formation of human adult stem cells from bone marrow stroma. *Cell Tissue Research*. 2005; 320(2):269-76.

- [53] Miljkovic ND, Cooper GM, Marra KG. Chondrogenesis bone morphogenetic protein-4 and mesenchymal stem cells. *Osteoarthritis Cartilage*. 2008; 16(10):1121-30.
- [54] Chubinskaya S, Hurtig M, Rueger DC. OP-1/BMP-7 in Cartilage repair. *International Orthopaedics*. 2007; 31(6):773-781.
- [55] An C, Cheng Y, Yuan Q, Li J. IGF-1 and BMP-2 induces differentiation of adipose-derived mesenchymal stem cells into chondrocytes-like cells. *Annals of Biomedical Engineering*. 2010; 38(4):1647-54.
- [56] O'Keeffe GW, Dockery P, Sullivan AM. Effects of growth/differentiation factor 5 on the survival and morphology of embryonic rat midbrain dopaminergic neurones in vitro. *Journal of Neurocytology*. 2004; 33(5):479-88.
- [57] Feng G, Wan Y, Balian G, Laurencin CT, Li X. Adenovirus-mediated expression of growth and differentiation factor-5 promotes chondrogenesis of adipose stem cells. *Growth Factors*. 2008; 26(3):132-42.
- [58] Kramer J, Hegert C, Guan K, Wobus AM, Muller PK, Rohwedel J. Embryonic stem cell-derived chondrogenic differentiation in vitro: activation by BMP-2 and BMP-4. *Mechanism of Development*. 2000; 92(2):193-205.
- [59] Schmidt MB, Chen EH, Lynch SE. A review of the effect of insulin-like growth factor and platelet derived growth factor on in vivo cartilage healing and repair. *Osteoarthritis Cartilage*. 2006; 14(5):403-12.
- [60] Kiep D, Ciarmatori S, Hoeflich A, Wolf E, Tonshoff B. Insulin-like growth factor (IGF)-I stimulates cell proliferation and induces IGF binding protein (IGFBP)-3 and IGFBP-5 gene expression in cultured growth plate chondrocytes via distinct signaling pathway. *Endocrine Society*. 2005; 146(7):3096-3104.
- [61] Longobardi L, O'Rear L, Aakula S, Johnstone B, Shimer K, Chytil A, et al. Effect of IGF-I in the chondrogenesis of bone marrow mesenchymal stem cells in the presence or absence of TGF- $\beta$  signaling. *Journal of Bone and Mineral Research*. 2006; 21(4):626-36.
- [62] Yammani RR, Loser RF. Extracellular nicotinamide phosphoribosyl transferase (NAMPT/Visfatin) inhibits insulin-like growth factor-1 signaling and proteoglycan synthesis in human articular chondrocytes. *Arthritis Research and Therapy*. 2012; 14(1):23.
- [63] Baxter RC. Insulin-Like growth factor (IGF)-binding protein: Interactions with IGFs and intrinsic bioactivities. *American Journal of Physiology-Endocrinology Metabolism*. 2000; 278(6):967-76.
- [64] Schneide MR, Wolf E, Hoeflich A, Lahm H. IGF-binding protein-5: Flexible player in the IGF system and effector on its own. *Journal of Endocrinology*. 2002; 172(3):423-40.
- [65] Kiepe D, Ciarmatori S, Hoeflich A, Wolf E, Tonshoff B. Insulin-Like Growth Factor (IGF)-I stimulates Cell Proliferation and Induces IGF Binding Protein(IGFBP)-3 and IGFBP-5 Gene Expression in cultured growth plate chondrocytes via distinct signaling pathways. *Journal of Endocrinology*. 2005; 146(7):3096-3104.
- [66] Starkman BG, Cravero JD, Delcarlo M, Loser RF. IGF-I stimulation of proteoglycan synthesis by chondrocytes requires activation of the PI 3-kinase pathway but not ERK MAPK. *Biochemical Journal*. 2005; 389(3):723-9.
- [67] Martin JA, Scherb MB, Lembke LA, Buckwalter J. Damage control mechanisms in articular cartilage: The role of the insulin-like growth factor I axis. *Iowa Orthopaedic Journal*. 2000; 20:1-10.
- [68] Headey SJ, Keizer DW, Yao S, Brasier G, Kantharidis P, Bach LA, et al. C-terminal domain of insulin-like growth factor (IGF) binding protein-6: structure and interaction with IGF-II. *Molecular Endocrinology*. 2004; 18(11):2740-50.
- [69] Itoh N, Ornitz DM. Fibroblast growth factors: from molecular evolution to roles in development, metabolism and disease. *Journal of Biochemistry*. 2011; 149(2):121-30.
- [70] Verkataraman G, Raman R, Sasisekharan V, Sasisekhar R. Molecular characteristics of fibroblast growth factor receptor heparin-like glycosaminoglycan complex. *Proceedings of the National Academy of Sciences of the United States of America*. 1999; 96(7):3658-63.
- [71] Kanda S, Miyata Y, Kanetake H, Smithgall TE. Fibroblast growth factor-2 induces the activation of Src through Fes, which regulates focal adhesion disassembly. *Experimental Cell Research*. 2006; 312(16):3015-22.
- [72] Stavridis MP, Lunn JS, Collins BJ, Storey KG. A discrete period of FGF-induced Erk1/2 signaling is required for vertebrate neural specification. *Development*. 2007; 134(16):2889-94.
- [73] Sørensen V, Zhen Y, Zakrzewska M, Haugsten EM, Wälchli S, Nilsen T, et al. Phosphorylation of fibroblast growth factor (FGF) receptor 1 at Ser777 by p38 mitogen-activated protein kinase regulates translocation of exogenous FGF1 to the cytosol and nucleus. *Molecular and Cellular Biology*. 2008; 28(12):4129-41.
- [74] Ohbayashi N, Shibayama M, Kurotaki Y, Imanishi M, Fujimori T, Itoh N, et al. FGF18 is required for normal cell proliferation and differentiation during osteogenesis and chondrogenesis. *Genes & Development*. 2002; 16(7):870-9.
- [75] Ellman MB, An HS, Muddasani P, Im HJ. Biological impact of the fibroblast growth factor family on articular cartilage and intervertebral disc homeostasis. *Gene*. 2008; 420(1):82-9.
- [76] Mandle EW, Jahr H, Koevoet JL, Vanlecuwen JP, Weinans H, Verhaar JA, et al. Fibroblast growth factor-2 in serum-free medium is a potent mitogen and reduces dedifferentiation of human ear chondrocytes in monolayer culture. *Matrix Biology*. 2004; 23(4):231-241.
- [77] Hashemibeni B, Zarei R, Esfandiari E, Valiani A. Adipose derived stem cells and application in musculoskeletal tissue repair. *Global Journal of Medicine Researches and Studies*. 2014; 1(1):11-16.
- [78] Rider DA, Dombrowski C, Sawyer AA, Ng GH, Leong D, Huttmacher DW, et al. Autocrine fibroblast growth factor 2 increases the multipotentiality of human adipose-derived mesenchymal stem cells. *Stem Cells*. 2008; 26(6):1598-608.
- [79] Yan D, Chen D, Cool SM, Van Wijnen AJ, Mikecz K, Murphy G, et al. Fibroblast growth factor receptor 1 is principally responsible for fibroblast growth factor 2-induced

- catabolic activities in human articular chondrocytes. *Arthritis Research and Therapy*. 2011; 13(4):130.
- [80] Moore EE, Bendele AM, Thompson DL, Littau A, Waggle KS, Reardon B, et al. Fibroblast growth factor-18 stimulates chondrogenesis and cartilage repair in a rat model of injury-induced osteoarthritis. *Osteoarthritis Cartilage*. 2005; 13(7):623-631.
- [81] Davidson D, Blance A, Filion D, Wang H, Plut P, Pfeffer G, Buschmann M.D, Henderson J.E. Fibroblast growth factor (FGF)18 signals through FGF receptor-3 to promote chondrogenesis. *Journal of Biological Chemistry*. 2005; 280(21):20509-515.
- [82] Vincent TL. Explaining the fibroblast growth factor paradox in osteoarthritis: Lessons from conditional knockout mice. *Arthritis & Rheumatism*. 2012; 64(12):3835-8.
- [83] Bessa PC, Casal M, Reis RL. Bone morphogenetic proteins in tissue engineering: The road from laboratory to clinic, part II (BMP delivery). *Journal of Tissue Engineering and Regenerative Medicine*. 2008; 2(2-3):81-96.
- [84] Fortier LA, Barker JU, Strauss EJ, McCarrel TM, Cole BJ. The role of growth factors in cartilage repair. *Clinical Orthopaedics and Related Research*. 2011; 469(10):2706-15.