

# Comparisons between Two Antioxidant Butylated Hydroxytoluene and Glutathione Supplemented Cryopreservation Medium on Human Sperm DNA Integrity


Marzieh Ghorbani <sup>1</sup>, Heidar Tavilani <sup>2</sup>, Iraj Khodadadi <sup>3,4</sup>, Iraj Amiri <sup>1\*</sup>

1. Endometrium and Endometriosis Research Center; Hamadan University of Medical Sciences, Hamadan, Iran.

2. Urology and Nephrology Research Center; Hamadan University of Medical Sciences, Hamadan, Iran.

3. Department of Biochemistry, Faculty of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran.

4. Urology and Nephrology Research Center; Hamadan University of Medical Sciences, Hamadan, Iran.



Iraj Amiri has PhD in the field of Anatomy from Tehran University of Medical Sciences and he is a member of the scientific board of Hamedan University of Medical Sciences. He is currently a full professor at the university and is the head of Department of Anatomy and director of Embryology laboratory at Fatemeh Infertility Center.

Article info:

Received: 04 Feb. 2015

Accepted: 21 Apr. 2015

## ABSTRACT

**Introduction:** Cryopreservation of semen is routinely used in a variety of circumstances including before assisted reproduction treatments, pre- radiation or chemotherapy treatment and etc. The aim of this study was to compare the effect of Butylated hydroxytoluene (BHT) and Glutathione supplemented cryopreservation medium on sperm parameters and amount of DNA fragmentation during the freeze-thaw process.

**Methods:** Semen samples were obtained from 60 donors. After the determination of basic parameters, groups of three sample with similar parameters were pooled and processed by Pure Sperm gradient centrifugation. The semen samples were then diluted with normal freezing medium (control) or a medium containing 5mM glutathione (test) and 0.5 mM BHT (test) stored in liquid nitrogen. Frozen cryovials were thawed individually for 20 seconds in a water bath (37 °C) for evaluation.

**Results:** Significant differences were observed in motility, viability and DNA fragmentation. Motility and viability were significantly higher in treated groups with 0.5 Mm in 5 min BHT than the control group and Glutathione 5mM (P<0.001).

**Conclusion:** Significant differences were observed in motility, viability and DNA fragmentation. Motility and viability were significantly higher in treated groups with 0.5 Mm in 5 min BHT than the control group and Glutathione 5mM (P<0.001).

### Key Words:

Butylated hydroxytoluene, Glutathione, Cryopreservation, DNA fragmentation, Spermatozoa

### \* Corresponding Author:

Iraj Amiri, PhD

Address: Endometrium and Endometriosis Research Center; Hamadan University of Medical Sciences, Hamadan, Iran.

Tel: +98 (912) 3362561

E-mail: amiri44@yahoo.com

## 1. Introduction

**C**ryopreservation is the procedure that stabilizes and preserve the cells at low temperatures and it is commonly used in infertility clinics and sperm banks. Cryopreservation was discovered in 1948 by polg-smith-parks to demonstrate that sperm of poultry and cattle can survive at 70 °C in the presence of glycerol, which was in fact the theory of freezing sex cells and tissues. Sperm freezing is routinely used for pre-radiation, chemotherapy treatment, for men undergoing vasectomy, cancer, storage of donor semen for HIV and Hepatitis patients. It is also used for Azoospermic patients who have done the practice testicular sperm extraction (TESE) or percutaneous epididymal sperm aspiration (PESA) [1]. Despite tremendous efforts, Cryopreservation leads to incomplete morphological, biochemical, chromosomal changes (formation of polyploidy), ionic disturbances and damages the membrane [1]. It is proved that the freezing process leads to reduced motility and viability and increased sperm DNA damage [2, 3].

Damage in sperm cell and membrane injury may be related to cold shock, intracellular ice crystal formation and osmotic stress during cryopreservation [4]. It seems that freezing and thawing gamete results in an increased amount of reactive oxygen species (ROS) which could damage cellular compounds including DNA, lipids, and proteins [2, 3].

Recently scientists have tried to reduce the effects of ROS and improve quality of thawed sperms with the addition of antioxidant supplements to the freezing medium. For this purpose, antioxidants such as ascorbate [5], vitamin E and reduced glutathione [6] have been used. Butylated hydroxytoluene (BHT), synthetic phenolic antioxidant with Lipophilic properties, could control peroxy radicals to hydro peroxides and reduce auto-oxidation. BHT has been tested in animal studies [7-9].

Aitken and Clarkson [1988] showed that the addition of BHT into the sperm freezing media decreases ROS production, thereby reducing damage to DNA. Another antioxidant which used is glutathione non-protein thiol with low molecular weight in plant and animal cells and its intracellular concentration 5-10 mmol while the extracellular one is third of this amount. Glutathione is a tripeptide which has gamma peptide linkage and therefore it is almost stable in the intracellular since the intracellular peptidases can cleave peptide bonds formed by the alpha-carboxyl groups of amino acids [6, 10].

The main objective of this study was to evaluate DNA fragmentation in sperm after thawing in the presence of BHT antioxidant and glutathione.

## 2. Materials and Methods

### Semen collection and processing

The study was approved by the Ethical Committee of Hamadan University of Medical Sciences and written consent was obtained from all participants. Semen samples were obtained from 60 donors with normal semen analysis by masturbation according to WHO [11] guidelines and 60 semen samples with similar characteristics were randomly divided into 20 sets of three semen samples.

Aliquots of 1ml of the liquefied pooled semen samples were layered on top of the upper layer of 40% and 80% Pure Sperm gradient (Nidacon International, Sweden) and centrifuged at 300 g for 20 minutes. After removal of the supernatant, the pellets were washed with 5ml of cook sperm medium (COOK IVF, Australia) and centrifuged at 500 g for 10 minutes [12].

Then, the pellets were resuspended in 0.5 ml of cook sperm medium. Finally, sperm samples were transferred into a series of cryotubes (5 tubes) containing 0.5 mM BHT concentrations and tubes were incubated for 5 minutes at 37 °C to allow uptake of BHT by spermatozoa. Then, semen samples were slowly frozen in liquid nitrogen vapors [7] and stored at -196 °C.

In the another group, 5 mM Glutathione was added to freezing medium. After 10 min equilibration of sperms at room temperature in this solution, the mixture are slowly frozen in liquid nitrogen vapors for 15 min and placed in special containers and they are stored in the (-196 °C liquid N<sub>2</sub>). In the last group (control group), sperms were freezed in freezing medium without addition of any antioxidant.

### Sperm thawing

Frozen sperm vials were placed in water bath (37 °C) for 4-5 minutes. Once the specimen were completely thawed, then gently were mixed and washed with sperm medium to eliminate cryosperm [7]. After thawing, sperm motility, viability (detected by staining with eosin 0.5%), and DNA fragmentation were evaluated.

### Viability test

The viability of the sperm in the sample was assessed by means of eosin 0.5% stain. The sperm smears were

**Table 1.** Level of sperm motility, viability and DNA fragmentation in three groups. Sample (n=20)

Group	Motility (%)	Viability (%)	Damaged DNA(%)
Control	34.7±5.7	46.56±5.9	49±7.6
BHT 0.5 mM/5 min	44.4±5.9	56.04±5.7	25.3±7.6
Glutathione 5 mM	35.39±5.24	49.56±8.51	36.13±8.21

ANATOMICAL SCIENCES

prepared by mixing a drop of semen with one drop of the stain on a warm slide. The viability was assessed by counting 200 cells under 400\* magnification. Sperm with partial or complete red colorization were assigned non-viable or dead whereas colorless sperm were designated as alive [11].

### Evaluation of sperm DNA fragmentation

Determination of sperm DNA fragmentation was carried out using Cell Death Detection Kit (Roche Diagnostics, Deutschland GmbH, Germany). Thawed sperm were smeared on microscope slides, air-dried, fixed with 4% paraformaldehyde in phosphate-buffered saline for 60 min and permeabilized with 0.1% Triton X-100 in 0.1% sodium citrate. Then, smears of sperm were further processed for TUNEL assay. Spermatozoa with fragmented DNA were visualized using fluorescence microscope in 488 nm (Figure A,B). A negative control (sperm without the addition of the Tdt enzyme) and a positive control (sperm treated with a 141 mM solution of H<sub>2</sub>O<sub>2</sub>) were also assessed by TUNEL assay [10].

Determination of DNA fragmentation in the thawed sperm of control group (A), no addition of Antioxidant) and in the thawed sperm with Antioxidants (B) using the TUNEL test. Spermatozoa with fragmented DNA were visualized using fluorescence microscope in excitation wavelength 488 nm and detection in the 540 nm. The spermatozoa with DNA fragmentation are shown in green and spermatozoa having intact DNA are shown in red.

### Statistical analysis

Results were expressed as the mean±SD. Sperm motility, motion characteristics motility, viability and DNA fragmentation were analyzed by analysis of variance, followed by Tukey's post hoc test to determine significant differences between groups. Differences with values of P<0.05 were considered to be statistically significant. Statistical analyses were performed by using SPSS 13 package program.

### 3. Results

Semen parameters of all participants are illustrated in Table 1. As shown in the table, significant differences were observed in motility, viability and DNA fragmentation between the groups. Motility were significantly higher in treated group with 0.5 Mm in 5 min BHT (44.4±5.9) than the control group (34.7±5.7) and Glutathione 0.5mM (35.39±5.24) (P<0.001). Viability were significantly higher in treated group with 0.5 Mm in 5 min BHT (56.04±5.7) than the control group (46.56 ±5.9) and Glutathione 0.5mM (49.56±8.51) (P<0.001). DNA fragmentation were significantly lower in treated groups (25.3±7.6) than the control (49±7.6) and Glutathione 0.5mM (36.13±8.21).

### 4. Discussion

The results of this study demonstrate that the addition of 0.5 mM BHT to the freezing medium decreased DNA fragmentation compared to the controls and Glutathione 0.5mM (P<0.001). Furthermore, freezing medium supplemented with 0.5 mM BHT led to higher sperm motility and viability compared with control Glutathione 0.5 mM (P<0.001).

Freezing and thawing process has deleterious effects on sperm DNA integrity by inducing DNA fragmentation [3]. DNA integrity of sperm is necessary for the correct transmission of genetic information to the next offspring and therefore, DNA damage may result in male infertility [13]. Studies have proved that high levels of ROS that are created during freezing and thawing process modify the occurrence of single- and double-strand DNA breakage in the sperm nucleus. Here, we showed that the addition of BHT into the freezing media reduces the percent of sperm with DNA damage from 49% in non-treated cells to 25% in BHT-received cells. Therefore, it can be concluded that the addition of BHT to the freezing media reduces spermatozoa ROS formation and DNA fragmentation. Tuncer et al., 2010 reported that adding Glutathione to the freezing medium have an impact on the DNA integrity and reduction of DNA damage [14-16]. Freezing and thawing process, leading to loss of DNA integrity and induces single- and double-strand DNA break of sperm cells. Our

result indicated that the addition of BHT(0.5 mM) to the sperm freezing media, significantly reduced sperm DNA damage compare to control and Glutathione groups.

We conclude that these antioxidants can be good scavenger of free radicals in the spermatozoa freezing medium and can compensate reduced activity of superoxide dismutase and glutathione peroxidase after a cycle of freezing and thawing. It seems that BHT could be an important scavenger of free radicals in the spermatozoa freezing medium and can compensate reduced activity of superoxide dismutase and glutathione peroxidase after a cycle of freezing and thawing. In conclusion, the addition of BHT to the freezing medium could be of advantage in reduction of ROS formation and prevents detrimental effect of ROS to human sperm function.

### Acknowledgement

This research was financially supported by Hamadan University of Medical Sciences, Hamedan, Iran.

### References

- [1] Donnelly ET, McClure N, Lewis SE. Cryopreservation of human semen and prepared sperm: effects on motility parameters and DNA integrity. *Fertility and Sterility*. 2001; 76(5):892-900.
- [2] Thomson LK, Fleming SD, Aitken RJ, De Iulius GN, Zieschang J-A, Clark AM. Cryopreservation-induced human sperm DNA damage is predominantly mediated by oxidative stress rather than apoptosis. *Human Reproduction*. 2009; 24(9):2061-70.
- [3] Zribi N, Chakroun NF, Ben Abdallah F, Elleuch H, Sellami A, Gargouri J, et al. Effect of freezing-thawing process and quercetin on human sperm survival and DNA integrity. *Cryobiology*. 2012; 65(3):326-31.
- [4] Watson P. The causes of reduced fertility with cryopreserved semen. *Animal Reproduction Science*. 2000; 60-61:481-92.
- [5] Li Z, Lin Q, Liu R, Xiao W, Liu W. Protective effects of ascorbate and catalase on human spermatozoa during cryopreservation. *Journal of Andrology*. 2010; 31(5):437-44.
- [6] Gadea J, Molla M, Selles E, Marco M, Garcia-Vazquez F, Gardon J. Reduced glutathione content in human sperm is decreased after cryopreservation: effect of the addition of reduced glutathione to the freezing and thawing extenders. *Cryobiology*. 2011; 62(1):40-6.
- [7] Ijaz A, Hussain A, Aleem M, Yousaf M, Rehman H. Butylated hydroxytoluene inclusion in semen extender improves the post-thawed semen quality of Nili-Ravi buffalo (*Bubalus bubalis*). *Theriogenology*. 2009; 71(8):1326-9.
- [8] Roca J, Gil MA, Hernandez M, Parrilla I, Vazquez JM, Martinez EA. Survival and Fertility of Boar Spermatozoa After Freeze-Thawing in Extender Supplemented With Butylated Hydroxytoluene. *Journal of Andrology*. 2004; 25(3):397-405.
- [9] Memon AA, Wahid H, Rosnina Y, Goh YM, Ebrahimi M, Nadia F. Effect of ascorbic acid concentrations, methods of cooling and freezing on Boer goat semen cryopreservation. *Reproduction in Domestic Animals*. 2013; 48(2):325-30.
- [10] Greco E, Romano S, Iacobelli M, Ferrero S, Baroni E, Minasi MG, et al. ICSI in cases of sperm DNA damage: beneficial effect of oral antioxidant treatment. *Human Reproduction*. 2005; 20(9):2590-4.
- [11] World Health Organization. WHO laboratory manual for the examination and processing of human semen. Geneva: World Health Organization Publication; 2010.
- [12] Amiri I, Ghorbani M, Heshmati S. Comparison of the DNA fragmentation and the sperm parameters after processing by the density gradient and the swim up methods. *Journal of Clinical and Diagnostic Research*. 2012; 6(9):1451.
- [13] Agarwal A, Said TM. Role of sperm chromatin abnormalities and DNA damage in male infertility. *Human Reproduction Update*. 2003; 9(4):331-45.
- [14] Gadea J, Molla M, Selles E, Marco MA, Garcia-Vazquez FA, Gardon JC. Reduced glutathione content in human sperm is decreased after cryopreservation: Effect of the addition of reduced glutathione to the freezing and thawing extenders. *Cryobiology*. 2011; 62(1):40-6.
- [15] Bucak MN, Sariözkan S, Tuncer PB, Sakin F, Ateşşahin A, Kulaksız R, et al. The effect of antioxidants on post-thawed Angora goat (*Capra hircus ancyrensis*) sperm parameters, lipid peroxidation and antioxidant activities. *Small Ruminant Research*. 2010; 89(1):24-30.
- [16] Tuncer PB, Bucak MN, Büyüklebici S, Sariözkan S, Yeni D, Eken A, et al. The effect of cysteine and glutathione on sperm and oxidative stress parameters of post-thawed bull semen. *Cryobiology*. 2010; 61(3):303-7.